

THE ZETA FUNCTIONS OF THE FANO SURFACES OF CUBIC THREEFOLDS

XAVIER ROULLEAU

ABSTRACT. We give an algorithm to compute the zeta function of the Fano surface of lines of a smooth cubic threefold $F \hookrightarrow \mathbb{P}^4$ defined over a finite field. We obtain some examples of Fano surfaces with supersingular reduction.

1. INTRODUCTION.

Let $k = \mathbb{F}_q$ be a finite field of characteristic $p \neq 2$ and let $F \hookrightarrow \mathbb{P}_{/k}^4$ be a smooth cubic hypersurface defined over k . Through a generic point of $\bar{F}$ pass 6 lines of $\bar{F}$. The variety that parametrizes the lines on F is a smooth projective surface defined over k called the Fano surface of lines of F . This surface S is minimal of general type.

In [14], we prove that the Tate conjecture holds for S . Recall that the zeta function $Z(X, T)$ of a smooth variety $X_{/k}$ encodes the number of rational points of X over the extensions $k_r = \mathbb{F}_{q^r}$. The Tate conjecture for X predicts that the Picard number ρ_X of $X_{/k}$ equals to the order of pole of the zeta function $Z(X, q^{-s})$ at $s = 1$.

Knowing that the Tate conjecture holds for a Fano surface S , it is then natural to study its consequences and applications : this is the aim of this paper. Using the results of Bombieri and Swinnerton-Dyer [5], we give an algorithm that compute the zeta function $Z(S, T)$ for a Fano surface S defined over k containing a k -rational point. We are therefore able to obtain the Picard number ρ_S , which is an important invariant but usually difficult to compute.

The intermediate Jacobian $J(F)$ of the cubic F is a certain 5-dimensional Abelian variety canonically associated to F . This variety $J(F)$ is isomorphic over k to the Albanese variety of S , and it is also a Prym variety $Pr(C/\Gamma)$ associated to a degree 2 cover $C \rightarrow \Gamma$ of a certain plane quintic curve Γ . As we will see, it is equivalent to compute the zeta functions $Z(F, T)$, $Z(S, T)$ or $Z(J(F), T)$.

In [6], Kedlaya gives an algorithm to compute the zeta function of cubic threefolds F that are triple abelian cover of $\mathbb{P}^3$ branched over a smooth cubic surface. This algorithm runs by computing the number of points on the cubic hypersurface $F \hookrightarrow \mathbb{P}^4$ and uses the order 3 symmetries of F in order to reduce the computations. The algorithm we give (implemented in Sage, see [13]) runs for any cubics containing a k -rational line. To our knowledge, it seems

also to be the first algorithm proposed in order to compute the zeta function of a Prym variety.

We then study some interesting examples of cubic threefolds and compute the zeta functions of their Fano surfaces over fields of characteristic 5 and 7. We study also the Fano surface $S/\mathbb{Q}$ of the Klein cubic threefold :

$$F/\mathbb{Q} = \{x_1^2x_2 + x_2^2x_3 + x_3^2x_4 + x_4^2x_5 + x_5^2x_1 = 0\}.$$

The surface S has good reduction S_p at every prime $p \neq 11$.

Corollary 1. *Let be $p \neq 2, 11$. If 11 is a square modulo p , then S_p has geometric Picard number 25, otherwise S_p is supersingular, i.e. its geometric Picard number equals $45 = b_2$.*

To the knowledge of the author, there are only a few cases where the zeta function of surfaces have been computed : apart from the cases of Abelian surfaces and a product of two curves, the zeta function has been computed for some examples of $K3$, some elliptic surfaces, for Fermat and some Delsarte hypersurfaces (see [17] and the references therein).

Acknowledgements. The author gratefully thanks Bert van Geemen for his suggestion to consider the conic bundle structure for computing the zeta function of cubics, and Kiran Kedlaya for emails exchanges. This paper has been supported under FCT grant SFRH/BPD/72719/2010 and project Géometria Algébrica PTDC/MAT/099275/2008.

2. THE ZETA FUNCTION OF A CUBIC THREEFOLD AND ITS FANO SURFACE.

2.1. Notations, hypothesis. Let F be a smooth cubic hypersurface defined over a field k of characteristic not 2 and let S be its Fano surface. The surface S is a smooth geometrically connected variety defined over k [2, Thm 1.16 i and (1.12)]. We will suppose that S has a rational point s_o i.e. that F contains a line defined over k . The Albanese variety A of S is 5 dimensional and is defined over k [1, Lem. 3.1]. Let $\vartheta : S \rightarrow A$ be the Albanese map such that $\vartheta_{s_o} = 0$. Let Θ be the (reduced) image of $S \times S$ under the map $(s_1, s_2) \rightarrow \vartheta_{s_1} - \vartheta_{s_2}$. The variety Θ is a divisor on A defined over k and (A, Θ) is a principally polarized abelian variety [4, Prop. 5]. In this paper, we call the pair $J(F) = (A, \Theta)$ the intermediate Jacobian of F .

2.2. Zeta functions, Weil polynomials. Let X be a smooth projective n -dimensional variety over a finite field $k = \mathbb{F}_q$ such that $\bar{X}$ (the variety over an algebraic closure $\bar{k}$ of k) is smooth. Let N_r be the number of rational points of $\bar{X}$ over $\mathbb{F}_{q^r}$. The zeta function of X is defined by $Z(X, T) = \exp(\sum_{r \geq 0} N_r \frac{T^r}{r})$ and can be written:

$$Z(X, T) = \sum_{i=0}^{i=2n} P_i(X, T)^{(-1)^{i+1}}$$

where $P_i(X, T)$ is the Weil polynomial with integer coefficients:

$$P_i(X, T) = \det(1 - \pi^* | H^i(\bar{X}, \mathbb{Q}_\ell)),$$

for π the Frobenius endomorphism of $\bar{X}$. Suppose that X is a surface. Let $Br(X)$ be the Brauer group of X and let $NS_k(X)$ be its Néron-Severi group.

Theorem 2. ([9], *Artin-Tate Conjecture*). *Suppose that the Tate conjecture is satisfied by X and the characteristic of k is at least 3. The Brauer group of X is finite and:*

$$\lim_{s \rightarrow 1} \frac{P_2(X, q^{-s})}{(1 - q^{1-s})^{\rho_X}} = \frac{(-1)^{\rho_X-1} |Br(X)| \text{Disc}(NS_k(X))}{q^{\alpha(X)} |NS_k(X)_{\text{tor}}|^2}$$

where $\alpha(X) = \chi(X, \mathcal{O}_X) - 1 + \dim \text{PicVar}(X)$, $NS_k(X)_{\text{tor}}$ is the torsion sub-group of $NS_k(X)$, ρ_X and $\text{Disc}(NS_k(X))$ are respectively the rank and the absolute value of the discriminant of $NS_k(X)/NS_k(X)_{\text{tor}}$.

Note that the order of a finite Brauer group is always a square. Let S be the Fano surface of a smooth cubic threefold $F \hookrightarrow \mathbb{P}^4$. The Picard variety $\text{PicVar}(S)$ of S is reduced of dimension 5 [20, Lem. 1.1], and $\chi = 6$, therefore $\alpha(S) = 10$.

As $H^2(\bar{S}, \mathbb{Q}_\ell) \simeq \wedge^2 H^1(\bar{A}, \mathbb{Q}_\ell) \simeq H^2(\bar{A}, \mathbb{Q}_\ell)$, the zeta function of S can be computed if we know the zeta function of A ; in fact we just need to know the action of the Frobenius on $H^1(\bar{A}, \mathbb{Q}_\ell)$.

Theorem 3. *Suppose characteristic $\neq 2$. The étale cohomology groups $H^3(\bar{F}, \mathbb{Q}_\ell)$ and $H^1(\bar{A}, \mathbb{Q}_\ell(1))$ are isomorphic as Galois modules.*

Proof. By [11, (14) and Thm 3], we have $H^3(\bar{F}, \mathbb{Q}_\ell) \simeq T_\ell(A) \otimes \mathbb{Q}_\ell(1)$ where $T_\ell(A)$ is the Tate module of A . But $T_\ell(A) \otimes \mathbb{Q}_\ell$ is isomorphic to $H^1(\bar{A}, \mathbb{Q}_\ell)$. $\square$

Remark 4. It would be interesting to know a bound on the order of the torsion subgroup $NS(S)_{\text{tor}}$. In the examples we give in Section 6, we compute the values of $\lim_{s \rightarrow 1} \frac{P_2(S, q^{-s})}{(1 - q^{1-s})^{\rho_S}}$. For all the limits we computed, we get results on the form $\frac{N}{q^{10}}$, $N \in \mathbb{Z}$, giving the idea that the group $NS(S)_{\text{tor}}$ should be trivial.

2.3. Zeta function of a smooth cubic threefold. The upper half part of the Hodge diamond of a smooth cubic threefold F over $\mathbb{C}$ is:

$$\begin{array}{ccccc} & & 1 & & \\ & & 0 & & 0 \\ & 0 & & 1 & & 0 \\ 0 & & 5 & & 5 & & 0 \end{array}$$

Thus $P_0(F, T) = 1 - T$, $P_1(F, T) = 1 = P_5(F, T)$, $P_2(F, T) = 1 - qT$. The polynomials P_4 and P_6 are computed by Poincaré duality: $P_4 = 1 - q^2T$, $P_6(F, T) = 1 - q^3T$.

Corollary 5. *We have:*

$$Z(F, T) = \frac{P_3(F, T)}{(1 - T)(1 - qT)(1 - q^2T)(1 - q^3T)},$$

and

$$N_r(F) = (1 + q^r + q^{2r} + q^{3r}) - q^r \text{Tr}(\pi^r | H^1(\bar{A}, \mathbb{Q}_\ell)),$$

where π is the Frobenius endomorphism of $\bar{A}$ and Tr denotes the trace.

The roots of $P_i(F, T)$ have absolute value $q^{-i/2}$. We have:

$$P_1(S, T) = P_1(A, T) = P_3(F, \frac{T}{q}).$$

Since by a computer it is possible to compute the number of points of F , it is possible in theory to get the zeta function of F, A and S . Kedlaya's algorithm computes the number of points of cubics F that are cyclic triple cover of $\mathbb{P}^3$. Alternatively, we can also compute the number of points of A , which is a Prym variety of a degree 2 étale cover of a certain plane quintic ; this is the algorithm we have implemented.

2.4. Zeta function and Picard number of a Fano surface. Let S be the Fano surface of a smooth cubic $F \hookrightarrow \mathbb{P}^4$ over $\mathbb{F}_q$. We have

$$Z(S, T) = \frac{P_1(S, T)P_3(S, T)}{(1 - T)P_2(S, T)(1 - q^2T)}.$$

We can write $P_1 = \prod_{i=1}^{i=10} (1 - \omega_i T) \in \mathbb{Z}[T]$ with $|\omega_i| = q^{-1/2}$. As

$$H^2(\bar{S}, \mathbb{Q}_\ell) = \wedge^2 H^1(\bar{S}, \mathbb{Q}_\ell)$$

we get $P_2(S, T) = \prod_{1 \leq i < j \leq 10} (1 - \omega_i \omega_j T)$. Moreover $P_3(S, T) = q^{15} T^{10} P_1(S, \frac{1}{q^2 T})$. Thus

$$(2.1) \quad Z(S, T) = \frac{\prod_{i=1}^{i=10} (1 - \omega_i T) \prod_{i=1}^{i=10} (1 - \frac{q^2}{\omega_i} T)}{(1 - T)(1 - q^2 T) \prod_{1 \leq i < j \leq 10} (1 - \omega_i \omega_j T)}.$$

The order of the pole at 1 of $Z(S, q^{-s})$ equals the number of elements of the set $\{(i, j) / 1 \leq i < j \leq 10 \text{ and } \omega_i \omega_j = q\}$; it is the multiplicity of the root $1/q$ in $P_2(S, T) = P_2(A, T)$. The Tate conjecture for Fano surfaces proved in [14] predicts that the Picard number ρ_S of S is equal to the order of pole of the zeta function $Z(S, q^{-s})$ at $s = 1$, thus by 2.1, we get:

Corollary 6. *The Fano surface has Picard number at least 5.*

Let $N_r(X)$ be the number of rational points of a variety X over $\mathbb{F}_{q^r}$. For each Weil polynomial $P_i(X, T)$, let $\omega_{i,1}, \dots, \omega_{i,b_i}$ be the reciprocal roots of P_i . The formula expressing the zeta function gives:

$$N_r(X) = \sum_{i=0}^{i=2 \dim X} (-1)^i \sum_{j=1}^{j=b_i} \omega_{i,j}^r.$$

We will use in section 4 this formula for computing the numbers $N_1(S), N_2(S)$ of the Fano surface S from the knowledge of the reciprocal roots $\omega_{1,1}, \dots, \omega_{1,10}$ of $P_1(S, T)$.

3. THE INTERMEDIATE JACOBIAN AS A PRYM VARIETY.

Let $F \hookrightarrow \mathbb{P}^4$ be a smooth cubic threefold defined over the finite field $k = \mathbb{F}_q$ of characteristic not 2. We will suppose that F contains a line L defined over k , or equivalently that the Fano surface contains a k -rational point.

The aim of this section is to give an algorithm that compute the zeta function of the cubic F , the Fano surface S and the Albanese variety A by using the conic bundle structure on an appropriate blow-up of F .

The algorithm will use two auxiliary curves : C_L the incidence curve parametrizing the lines on F meeting L and Γ_L the curve parametrizing planes Y containing L and such that the degree 3 plane curve $Y \cap F$ is the union of three lines.

Let $x_1, \dots, x_5$ be coordinates in $\mathbb{P}^4$ and let $L \hookrightarrow F$ be a k -rational line on F . We can suppose that $L = \{x_1 = x_2 = x_3 = 0\}$ and $F = \{F_{eq} = 0\}$ with

$$F_{eq} = \ell_1 x_4^2 + 2\ell_2 x_4 x_5 + \ell_3 x_5^2 + 2q_1 x_4 + 2q_2 x_5 + f,$$

where ℓ_1, ℓ_2, ℓ_3 are linear, q_1, q_2 are quadratic and f is a cubic homogenous forms in the variables x_1, x_2, x_3 . Let $X \hookrightarrow \mathbb{P}^4$ be the plane $x_4 = x_5 = 0$. Any plane $Y \hookrightarrow \mathbb{P}^4$ containing the line L meets X into a unique point. Thus the plane X parametrizes planes in $\mathbb{P}^4$ containing L . For a $\bar{k}$ -point $x = (x_1 : x_2 : x_3 : 0 : 0)$ of $X = \mathbb{P}^2$, we denote by Y_x the unique plane containing x . The intersection of F and Y_x has equation $F_{eq}(y_1 x_1 : y_1 x_2 : y_1 x_3 : y_2 : y_3) = 0$, i.e.:

$$y_1(\ell_1 y_2^2 + 2\ell_2 y_2 y_3 + \ell_3 y_3^2 + 2q_1 y_1 y_2 + 2q_2 y_1 y_3 + y_1^2 f) = 0$$

in the plane Y_x with coordinates y_1, y_2, y_3 . Therefore, the intersection $Y \cap F$ equals $L + Q_x$ where

$$Q_x = \{\ell_1 y_2^2 + 2\ell_2 y_2 y_3 + \ell_3 y_3^2 + 2q_1 y_1 y_2 + 2q_2 y_1 y_3 + y_1^2 f = 0\}.$$

The conic Q_x is irreducible over $\bar{k}$ if and only if it is smooth. The scheme parametrizing the planes Y_x such that Q_x decomposes (over an extension) as $Q_x = L_1 + L_2$ with L_1, L_2 two lines, is a plane reduced quintic curve $\Gamma_L \hookrightarrow X = \mathbb{P}^2(x_1, x_2, x_3)$ defined over k , of equation $\det M_\Theta = 0$, where:

$$M_\Theta = \begin{pmatrix} \ell_1 & \ell_2 & q_1 \\ \ell_2 & \ell_3 & q_2 \\ q_1 & q_2 & f \end{pmatrix}.$$

Let C_L be the incidence scheme parametrizing the lines that meet L .

Lemma 7. ([5, Lem. 2, 4], [12, Pro. (1.25)]). *The plane quintic curve Γ_L has at most nodal singularities. A point x on Γ_L is singular if and only if $Y_x \cap F = L + 2L'$, where L' is a line. The curve Γ_L is smooth for L generic. The scheme C_L is a reduced one dimensional subscheme of the Fano surface S . There is a natural degree 2 map $\mu : C_L \rightarrow \Gamma_L$ that is ramified precisely over the singularities of Γ . The point p of C_L is smooth if and only if $\mu(p)$ is smooth.*

If Γ_L is smooth, the Albanese variety of S is isomorphic to the Prym variety $Pr(C_L/\Gamma_L)$.

Let $\tilde{F} \rightarrow F$ be the blow-up of F along a line L on F . The variety $\tilde{F}$ has a natural structure of a conic bundle:

$$\tilde{F} \rightarrow \mathbb{P}^2 = X$$

where X parametrizes the planes containing the line L . The fiber over the point x is (isomorphic to) the quadric Q_x such that $F \cap Y_x = Q + L$. For x defined over k , the number of k -points of Q_x is:

- i) $q + 1$ if Q_x is geometrically irreducible i.e. $x \notin \Gamma_L(k)$ or if $x \in \Gamma_L(k)$ is a singular point of Γ , in that case $Q_x = 2L'$ with L' a k -rational line on F .
- ii) $2q + 1$ if $Q_x = L_1 + L_2$ with L_1, L_2 two different lines defined over k , i.e. the 2 points in C_L over $x \in \Gamma_L(k)$ are in $C_L(k)$.
- iii) 1 if the conic Q_x degenerates to a union of two line over a (degree 2) extension of k , i.e. if the 2 points in C_L over $x \in \Gamma_L(k)$ are not in $C_L(k)$.

Let $N_r(X)$ denotes the number of rational points of a variety X over the degree r extension $k_r = \mathbb{F}_{q^r}$ of k . The following Proposition is [5, Formula

(18)], however we reproduce the proof here because it explains the algorithm computing $N_r(F)$ we describe below.

Proposition 8. *We have*

$$N_r(F) = q^{3r} + q^{2r} + q^r + 1 + q^r(N_r(C_L) - N_r(\Gamma_L)).$$

Proof. Let a be the number of $\mathbb{F}_q$ -rational singularities of Γ_L (and C_L). Taking care of the three above possibilities i), ii) and iii), we get:

$$N_1(\tilde{F}) = (q+1)(N_1(\mathbb{P}^2) - N_1(\Gamma_L) + a) + (2q+1)\frac{1}{2}(N_1(C_L) - a) + (N_1(\Gamma_L) - a - \frac{1}{2}(N_1(C_L) - a))$$

thus

$$N_1(\tilde{F}) = q^3 + 2q^2 + 2q + 1 + q(N_1(C_L) - N_1(\Gamma_L)).$$

As each point on $L \hookrightarrow F$ is replaced by a $\mathbb{P}^1$ on $\tilde{F}$, we have moreover:

$$N_1(\tilde{F}) = N_1(F) - (q+1) + (q+1)^2,$$

thus $N_1(F) = q^3 + q^2 + q + 1 + q(N_1(C_L) - N_1(\Gamma_L))$. $\square$

Let $\mu : C_L \rightarrow \Gamma_L$ be the degree 2 map and let x be a k -rational smooth point on Γ . In order to compute the numbers $N_r(C_L) - N_r(\Gamma_L)$, we need to understand when the two $\bar{k}$ -rational points in $\mu^{-1}x$ are k -rational. For $1 \leq i \leq 3$, let $\delta_i \in H^0(\Gamma_L, \mathcal{O}(a))$, $a = 2$ or 4 be the (i, i) -minor of the matrix M_Θ .

Proposition 9. *Let x be smooth point of Γ . There exists an integer $1 \leq i = i(x) \leq 3$ such that $\delta_i(x) \neq 0$. The curve C_L has two rational points over $x \in \Gamma_L(k)$ if and only if $-\delta_i(x) \in (k^*)^2$.*

Proof. Let x a k -rational smooth point of Γ and let $Q = Q_x$ such that $F \cap Y_x = L + Q$. The line $L = \{y_1 = 0\}$ meets $Q_x \hookrightarrow Y_x$ in the points such that:

$$y_1 = \ell_1 y_2^2 + 2\ell_2 y_2 y_3 + \ell_3 y_3^2 = 0.$$

Therefore, if $-\delta_3(x) = \ell_2^2(x) - \ell_1(x)\ell_2(x)$ is nonzero, the curve C_L has two rational points over $x \in \Gamma_L(k)$ if and only if $-\delta_3(x) \in (k^*)^2$.

For $\delta_3(x) = 0$, we only sketch the proof ; see also [3, Lemme 1.6] and its proof. In that case, we have $Y_x \cdot F = L + Q_x = L + L_1 + L_2$ with L_1, L_2 defined over $\bar{k}$ and meeting in a k -rational point p of L . It is possible to explicitly compute a model in $\mathbb{P}^3$ of (the degree 6) scheme $\mathcal{X}_p$ of lines in the cubic F going through p . By knowing $\mathcal{X}_p$, we can determine whether the two points on $\mathcal{X}_p$ corresponding to the lines L_1, L_2 are k -rational or not and this is so if and only if $-\delta_1$ is a nonzero square or $-\delta_2$ is a nonzero square. $\square$

Let us describe the algorithm for the computation of the numbers $N_r(F)$ and $N_r(C_L) - N_r(\Gamma_L)$.

The input data is a cubic threefold F over $\mathbb{F}_q$ containing a $\mathbb{F}_q$ -rational line L . To this data is associated the matrix M_Θ defined above whose determinant is the equation of the quintic $\Gamma_L \hookrightarrow X = \mathbb{P}^2$ (maybe singular). Then we compute $N_r = N_r(C_L) - N_r(\Gamma_L)$ as follows:

Initiate $N_r := 0$. For $x \in \mathbb{P}^2(\mathbb{F}_{q^r})$, if $\det(M_\Theta)(x) = 0$ then if $-\delta_3(x) \in (\mathbb{F}_{q^r}^*)^2$ then $N_r := N_r + 1$, else if $-\delta_3(x) \neq 0$, then $N_r := N_r - 1$, otherwise if $-\delta_1(x) \in (\mathbb{F}_{q^r}^*)^2$ then $N_r := N_r + 1$ else if $-\delta_1(x) \neq 0$, then $N_r := N_r - 1$, otherwise if $-\delta_2(x) \in (\mathbb{F}_{q^r}^*)^2$ then $N_r := N_r + 1$ else if $-\delta_2(x) \neq 0$ then $N_r := N_r - 1$ end if, end for.

The output N_r equals $N_r(C_L) - N_r(\Gamma_L)$. Remark that the $-\delta_i$ are transition functions of an invertible sheaf $\mathcal{L}$ on Γ_L such that $\mathcal{L}^{\otimes 2} = \omega_{\Gamma}$. The data of $\mathcal{L}$ corresponds to the degree 2 cover $\mu : C_L \rightarrow \Gamma_L$ and a point on Γ_L is singular if and only if $\forall 1 \leq i \leq 3, \delta_i(x) = 0$.

The knowledge of $N_1, \dots, N_5$ is enough to get the 5 first coefficients of the degree 10 polynomial $P_1(Pr(C_L/\Gamma_L), T) \in \mathbb{Z}[T]$ and the remaining 5 ones are determined by the symmetries of P_1 .

Remark 10. This algorithm for computing the action of the Frobenius on the Prym variety $Pr(C_L/\Gamma_L)$ is generalizable to other plane curves occurring as discriminant locus of other quadric bundles, see e.g. [3].

4. EXAMPLES

4.1. Reduction in characteristic 5 and 7 of a cubic threefold. Let $F \hookrightarrow \mathbb{P}^4$ be the cubic threefold with equation:

$$F_{eq} = x_1x_4^2 + 2x_2x_4x_5 + x_3x_5^2 + 2q_1x_4 + 2q_2x_5 + f,$$

where:

$$\begin{aligned} q_1 &= x_1^2 + 2x_2^2 + x_2x_3 + x_3^2 \\ q_2 &= x_1x_2 + 4x_2x_3 + x_3^2 \\ f &= x_2^2x_3 - (x_1^3 + 4x_1x_2^2 + 2x_2^3). \end{aligned}$$

The cubic F is smooth in characteristic 5, 7, 11 and 13 ; it is singular in characteristic 2, 3. The associated quintic curve $\Gamma = \Gamma_L$ is smooth in characteristic 3, 7, 11, 13, but singular in characteristic 2, 5.

Remark 11. The cubic $F/\mathbb{Q}$ and its Fano surface $S/\mathbb{Q}$ have bad reduction at the place 3, however the intermediate Jacobian $J(F) \simeq Pr(C_L/\Gamma_L)$ has good reduction. This is the same phenomena as for the curves and their Jacobian. We remark also that the curves Γ_L and C_L both have bad reduction at the place 5, but the associated Prym variety has good reduction.

We have implemented the algorithm in Sage. Using a personal laptop, it takes 5 minutes to obtain $P_1(S, T)$ for S over $\mathbb{F}_7$.

Over $\mathbb{F}_5$, we get (see [13]) :

$$P_1(S/\mathbb{F}_5, T) = (5T^2 + 1)(625T^8 + 50T^6 + 40T^5 - 6T^4 + 8T^3 + 2T^2 + 1).$$

The Fano surface $S/\mathbb{F}_5$ has Picard number 5 and contains 33 $\mathbb{F}_3$ -rational points. We have:

$$A_5 := \lim_{s \rightarrow 1} \frac{P_2(S/\mathbb{F}_5, 5^{-s})}{(1 - 5^{1-s})^5} = \frac{2^{18} \cdot 3^5 \cdot 157}{5^{10}}.$$

Over $\mathbb{F}_7$, we get:

$$\begin{aligned} P_1(S/\mathbb{F}_7, T) = & 1 + 4T + 15T^2 + 46T^3 + 159T^4 + 460T^5 + 1113T^6 \\ & + 2254T^7 + 5145T^8 + 9604T^9 + 16807T^{10}. \end{aligned}$$

It is an irreducible polynomial over $\mathbb{Q}$, therefore by the Honda-Tate Theorem [10, Theorems 2-3, App. I], the intermediate jacobian $J(F)$ of F is simple. The Fano surface $S/\mathbb{F}_7$ has Picard number 5 and 97 $\mathbb{F}_7$ -rational points. We obtain:

$$A_7 := \lim_{s \rightarrow 1} \frac{P_2(S/\mathbb{F}_7, 7^{-s})}{(1 - 7^{1-s})^5} = \frac{2^4 \cdot 83^2 \cdot 557 \cdot 5737}{7^{10}}.$$

Remark 12. We prove in [15] that a generic Fano surface over $\mathbb{C}$ has Picard number $\rho = 1$. One would like to exhibit an example of a Fano surface over $\mathbb{Q}$ with $\rho = 1$. By reducing the above Fano surface $S/\mathbb{Q}$ modulo a prime we obtain the bound $\rho_S \leq 5$. Since A_5/A_7 is not a square in $\mathbb{Q}$, we can apply the van Luijk method [8] and obtain the inequality $\rho_S \leq 4$.

Over the field $\mathbb{F}_{11}$, the computation becomes difficult : it needs 4 minutes to get N_4 but more that 24 hours to get N_5 . By [6, Lem. 1.2.3], for positive integers q, d, j and complex numbers $a_1, \dots, a_{j-1}$, there exists a certain disk of radius $\frac{d}{j}q^{j/2}$ which contains every a_j for which we can choose $a_{j+1}, \dots, a_d \in \mathbb{C}$ so that the polynomial

$$R(T) = 1 + \sum_{j=1}^{j=d} a_j T^j$$

has all roots on the circle $|T| = q^{-1/2}$. In our case, $\frac{d}{j}q^{j/2} = \frac{10}{5}11^{5/2} = 802.623\dots$ and we obtain that $P_1(S/\mathbb{F}_{11}, T) = Q_a$, where a is an integer in $\{80, \dots, 332\}$ and

$$Q_a = 1 - T + 13T^2 + T^3 - 28T^4 + aT^5 - 11 \cdot 28T^6 + 11^2T^7 + 11^3 \cdot 13T^8 - 11^4T^9 + 11^5T^{10}.$$

For all the consecutive values $a \in \{80, \dots, 332\}$, the polynomial Q_a has its roots equal to $11^{-1/2}$ (with error at most 10^{-10}) and we cannot distinguish the a corresponding to our Fano surface S .

4.2. The Klein cubic threefold. Let $S/\mathbb{Q}$ be the Fano surface of lines of the Klein cubic threefold :

$$F/\mathbb{Q} = \{x_1^2x_2 + x_2^2x_3 + x_3^2x_4 + x_4^2x_5 + x_5^2x_1 = 0\}.$$

It is easy to check that F (hence S) has good reduction at every prime $p \neq 11$.

Proposition 13. *Let us suppose $p \neq 2$. If 11 is not a square modulo p , then S_p the reduction mod p is supersingular i.e. its geometric Picard number equals $45 = b_2$, otherwise S_p has geometric Picard number 25.*

Proof. Let be $\nu = \frac{-1+\sqrt{-11}}{2}$, $\mathcal{O} = \mathbb{Z}[\nu]$ and $E = \mathbb{C}/\mathbb{Z}[\nu]$. The intermediate jacobian $J(F)_{/\mathbb{C}}$ is isomorphic to E^5 (see [16]). By [18, App. A3], the elliptic curve E has the following model over $\mathbb{Q}$:

$$y^2 + y = x^3 - x^2 - 7x + 10,$$

which we still denote by E . The curve E has good reduction for prime $p \neq 11$ and it has complex multiplication by $\mathcal{O}$ (over a certain extension). We use the criteria of Deuring [7, Chap. 13, Thm 12)] : for odd $p \neq 11$, the reduction of E modulo p is a supersingular if and only if p is inert or ramified in $\mathcal{O}$. By classical results on number theory, an odd prime $p \neq 11$ is inert or ramified in $\mathcal{O}$ if and only if 11 is not a square modulo p .

Over an extension, the intermediate Jacobian $J(F)$ is isogenous to E^5 . By [10], the geometric Picard number of the reduction modulo p of $J(F)$ is therefore 45 if 11 is not a square modulo p , and 25 otherwise. $\square$

RÉFÉRENCES

- [1] Achter J., "Arithmetic Torelli maps for cubic surfaces and threefolds", preprint arXiv :1005.2131v3
- [2] Altman A., Kleiman S., "Foundations of the theory of Fano schemes", *Compositio Math.* 34 (1977), no. 1, 3–47.
- [3] Beauville A., "Variétés de Prym et Jacobiennes intermédiaires", *Ann. Scient. Éc. Norm. Sup.* 10 (1977), 309–391.
- [4] Beauville A., "Les singularités du diviseur Θ de la jacobienne intermédiaire de l'hyper-surface cubique dans $\mathbb{P}^4$ ", *Algebraic threefolds (Varenna, 1981)*, pp. 190–208, *Lecture Notes in Math.*, 947, Springer, Berlin-New York, 1982.
- [5] Bombieri E., Swinnerton-Dyer H.P.F., "On the local zeta function of a cubic threefold", *Ann. Scuola Norm. Sup. Pisa* (3) 21 1967, 1–29.
- [6] Kedlaya K., "Effective p -adic cohomology for cyclic cubic threefolds", *Computational Algebraic and Analytic Geometry*, 127–171, *Contemp. Math.*, 572, Amer. Math. Soc., Providence, RI, 2012.
- [7] Lang S., "Elliptic Functions", Addison-Wesley Publishing Co., Inc., Reading, Mass.-London-Amsterdam, 1973. xii+326 pp.
- [8] van Luijk R., "K3 surfaces with Picard number one and infinitely many rational points", *Algebra and Number Theory*, Vol. 1, No. 1 (2007), 1–15.
- [9] Milne J., "On a conjecture of Artin and Tate", *Ann. of Math.* (2) 102 (1975), no. 3, 517–533.
- [10] Mumford D., "Abelian Varieties", *Tata Institute of Fundamental Research Studies in Mathematics*, No. 5 Published for the Tata Institute of Fundamental Research, Bombay; Oxford University Press, London, 1970.
- [11] Murre J.P., "Some results on cubic threefolds", *Classification of algebraic varieties and compact complex manifolds*, pp. 140–160. *Lecture Notes in Math.*, Vol. 412, Springer, Berlin, 1974.
- [12] Murre J.P., "Algebraic equivalence modulo rational equivalence on a cubic threefold", *Compositio Math.* 25 (1972), 161–206.
- [13] Roulleau X., Sage computations available in [http ://www-math.sp2mi.univ-poitiers.fr/~roulleau](http://www-math.sp2mi.univ-poitiers.fr/~roulleau/publications) (publications).
- [14] Roulleau X. "On the Tate conjecture for the Fano surfaces of cubic threefolds", *J. of Number Theory* 133 (2013), pp. 2320–2323.
- [15] Roulleau X., "Fano surfaces with 12 or 30 elliptic curves", *Michigan Math. J.* 60 (2011), no. 2, 313–329
- [16] Roulleau X., "The Fano surface of the Klein cubic", *J. Math. Kyoto Univ.* 49 (2009), no. 1, 113–129.
- [17] Schütt M., "K3 surfaces with non-symplectic automorphisms of 2-power order", *J. Algebra* 323 (2010), no. 1, 206–223.
- [18] Silverman J. "Advanced topics in the arithmetic of Elliptic curves", *Graduate Texts in Mathematics*, 151. Springer-Verlag, New York, 1994.
- [19] Tate J., "Algebraic cycles and poles of zeta functions", 965 *Arithmetical Algebraic Geometry (Proc. Conf. Purdue Univ., 1963)* pp. 93–110 Harper and Row, New York
- [20] Tyurin, A.N., "The geometry of the Fano surface of a nonsingular cubic $F \subset \mathbb{P}^4$ and Torelli Theorems for Fano surfaces and cubics", *Math. USSR Izv.* 5, 517–546 (1971)

Xavier Roulleau,
 Université de Poitiers,
 Laboratoire de Mathématiques et Applications, UMR 7348 du CNRS,
 Boulevard Pierre et Marie Curie,
 Téléport 2 - BP 30179,
 86962 Futuroscope Chasseneuil,
 France
 Xavier.Roulleau@math.univ-poitiers.fr